



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 39/155, 39/39</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/11019</b> <b>(43) International Publication Date:</b> 18 April 1996 (18.04.96)
<b>(21) International Application Number:</b> PCT/US95/12656 <b>(22) International Filing Date:</b> 2 October 1995 (02.10.95) <b>(30) Priority Data:</b> 08/318,480      5 October 1994 (05.10.94)      US <b>(71) Applicant:</b> VANDERBILT UNIVERSITY [US/US]; 405 Kirkland Hall, Nashville, TN 37240 (US). <b>(72) Inventors:</b> GRAHAM, Barney, S.; 2414 Barton Avenue, Nashville, TN 37212 (US). TANG, Yi-Wei; 2707 Barton Avenue, Nashville, TN 37212 (US). <b>(74) Agents:</b> BROOK, David, E. et al.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US).	<b>(81) Designated States:</b> AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.          Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

**(54) Title:** INTERLEUKIN-12 AS AN ADJUVANT FOR PARAMYXOVIRIDAE VACCINES**(57) Abstract**

A method is disclosed of reducing viral replication of a virus of the Paramyxoviridae family in a host, comprising administering to the host an antigen of the virus in combination with an effective adjuvant amount of interleukin-12 (IL-12). Human viruses of the Paramyxoviridae family include paramyxoviruses (e.g., parainfluenza virus 1, parainfluenza virus 2, parainfluenza virus 3 and parainfluenza virus 4), morbilliviruses (e.g., measles virus) and pneumoviruses (e.g., respiratory syncytial virus); other non-human viruses of the Paramyxoviridae family include canine distemper virus, bovine respiratory syncytial virus, Newcastle disease virus and rinderpest virus. A composition is also disclosed comprising a mixture of an antigen of a virus of the Paramyxoviridae family and an effective adjuvant amount of interleukin-12 (IL-12).

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

INTERLEUKIN-12 AS AN ADJUVANT FOR PARAMYXOVIRIDAE VACCINESBACKGROUND

Respiratory syncytial virus (RSV), a member of the *Pneumovirus* genus of the *Paramyxoviridae* family, is an important cause of respiratory disease in infants and children (Connors, M., et al., *J. of Virol.*, 66:7444-7451 (1992). The immunological basis for the differing susceptibility among individuals, and for the limited age range at which severe illness occurs, remains unclear.

The major impediment to advancing new candidate vaccines directed against RSV to clinical trials is an incomplete understanding of the vaccine-enhanced illness caused by formalin-inactivated RSV vaccines in the 1960's. Clinical trials of a formalin-activated alum-precipitated RSV vaccine in the 1960's showed that the vaccine elicited complement-binding antibodies but failed to protect against infection in children. In addition, the illness after subsequent infection was unusually severe with some deaths, and an increased rate of hospitalization (Kapikian, A.Z., et al. *Amer. J. Epidemiol.*, 89:405 (1969); Fulginiti, V.A., et al, *Amer. J. Epidemiol.*, 89:435 (1969); Kim, W.H., *Amer. J. Epidemiol.*, 89:422 (1969); Chin, J.R., et al. *Amer. J. Epidemiol.*, 89:449 (1969)). A similar enhanced illness can be induced in mice previously immunized with the formalin-inactivated vaccine upon RSV infection, but not in mice immunized with live RSV (Connors, M., et al., *J. Virol.* 66:7441 (1992); Graham, B.S., et al., *Immunol.* 151:2032 (1993); Alwan, W.H., et al., *J. Exp. Med.* 179:81 (1994)). Clinical trials of live attenuated RSV vaccine products have not been associated with enhanced illness. Although the live RSV vaccines did not result in enhanced pulmonary disease upon natural infection, the vaccines were, in other respects, as equally unsuccessful as the formalin-

-2-

- inactivated alum-precipitated RSV vaccines (Kim, W.H., et al., *Pediatrics*, 48:745 (1971); Kim, W.H., et al., *Pediatrics*, 52:56 (1973); Belshe, R.B., et al., *J. Infect. Dis.*, 145:311 (1982); Wright, R.B., et al., *Infect. Immun.*, 37:397 (1982). Temperature-sensitive mutants of RSV, cold-adapted RSV or live RSV given parenterally have been considered unsuccessful as vaccines because of high rates of reversion to wild-type, unacceptable virulence or lack of immunogenicity in the appropriate age group
- 10 (Graham, B.S., et al., *J. of Immun.*, 151:2032-2040 (1993). Thus, a need exists for development of efficacious methods of vaccination against RSV and for vaccine compositions.

#### SUMMARY OF THE INVENTION

- 15 The present invention is based on the discovery that IL-12 has a potent adjuvant effect for immunizing against Paramyxoviridae virus infection. In one embodiment, the invention comprises a method of reducing viral replication of a virus of the paramyxoviridae family in a host (e.g.,
- 20 mammalian, including human, and avian) comprising administering to the host an antigen of the virus in combination with an effective adjuvant amount of interleukin-12 (IL-12). Human viruses of the Paramyxoviridae family include paramyxoviruses (e.g.,
- 25 parainfluenza virus 1, parainfluenza virus 2, parainfluenza virus 3, parainfluenza virus 4 and mumps virus), morbilliviruses (e.g., measles virus) and pneumoviruses (e.g., respiratory syncytial virus); other non-human viruses of the Paramyxoviridae family include canine
- 30 distemper virus, bovine RSV, Newcastle disease virus and rhinderpest virus. In one embodiment, the invention relates to a method of reducing replication of the respiratory syncytial virus (RSV) in a host comprising administering to the host an antigen of RSV in combination

-3-

with an effective adjuvant amount of IL-12. Thus, the present invention also relates to a method of eliciting an immune response against viruses of the Paramyxoviridae family in a host, comprising administering to the host an antigen of a virus of the Paramyxoviridae family in combination with an effective adjuvant amount of IL-12. The present invention also relates to a method of immunizing a host against RSV comprising administering to the host a mixture comprising an antigen of respiratory syncytial virus in combination with an effective adjuvant amount of interleukin-12.

In addition, the present invention relates to a composition comprising a mixture of an antigen of a virus of the Paramyxoviridae family and an effective adjuvant amount of interleukin-12 (IL-12). In one embodiment, the invention relates to a composition comprising an antigen of the respiratory syncytial virus and IL-12.

As shown herein, exogenous IL-12 treatment administered at the time of immunization with RSV antigen, diminishes RSV replication and increases endogenous IL-12 mRNA expression at the time of subsequent RSV challenge. This results in a shift from a Th2 to a Th1-like pattern of cytokine expression and a consequent shift in antibody isotype utilization. These results demonstrate that IL-12 is a potent adjuvant for Paramyxoviridae vaccines.

#### DESCRIPTION OF THE FIGURES

Figure 1 is a bar graph of IL-12 treatment versus log 10 plaque forming units (pfu)/ gram lung from the plaque assay illustrating that IL-12 administered at the time of immunization has a marked effect on the reduction of viral replication.

Figure 2 is a bar graph of IL-12 treatment versus averaged  $\alpha$ -tubulin-normalized density from the mRNA Northern blots illustrating that IL-12 enhanced Th1 cell

-4-

differentiation and produced a shift from a Th2 to a Th1-like response in mice immunized with inactivated RSV immunogen.

#### DETAILED DESCRIPTION OF THE INVENTION

5       The present invention relates to a method of reducing viral replication of a virus of the Paramyxoviridae family in a host (e.g., mammal, including human, avian) comprising, administering to the host, a mixture of an antigen of the virus and an effective adjuvant amount of  
10 interleukin-12 (IL-12). Although the method of the present invention is exemplified using RSV, the method can be used to reduce viral replication of a variety of viruses from the Paramyxoviridae family which include human paramyxoviridae viruses, such as paramyxoviruses (e.g.,  
15 parainfluenza virus 1, parainfluenza virus 2, parainfluenza virus 3, parainfluenza virus 4 and mumps virus), morbilliviruses (e.g., measles virus) and pneumoviruses (e.g., respiratory syncytial virus). Other non-human viruses of the Paramyxoviridae family include canine  
20 distemper virus, bovine RSV, Newcastle disease virus and rhinderpest virus. In one embodiment, the method of the present invention is used to reduce viral replication of respiratory syncytial virus (RSV) in a host, and comprises administering to the host an RSV antigen and an effective  
25 adjuvant amount of IL-12.

In another embodiment, the method of the present invention is used to elicit an immune response against a virus of the Paramyxoviridae family in a host, comprising administering to the host an antigen of the virus and an  
30 effective adjuvant amount of IL-12. In addition, the present invention relates to a method of immunizing a host against RSV comprising administering to the host a mixture comprising an RSV antigen and an effective adjuvant amount of IL-12.

-5-

An antigen of a virus of the Paramyxoviridae family includes use of the whole virus (e.g., inactivated or live, attenuated whole virus), an antigenic portion of the virus, and recombinantly produced virus or portions thereof or  
5 fusion proteins. In addition, antigens of the present invention include nucleic acid sequences which encode an antigen of a virus of the Paramyxoviridae family. Antigenic portions of the viruses of the Paramyxoviridae family include the fusion glycoprotein (F protein) and the  
10 hemagglutinin-neuraminidase of the parainfluenza viruses 1, 2, 3, 4; the F protein and the hemagglutinin-neuraminidase of the mumps virus; the F protein and the hemagglutinin-neuraminidase of the measles virus; and the F protein and the G glycoprotein of the RSV. Other antigenic portions of  
15 the Paramyxoviridae family of viruses which can be used in the methods and compositions of the present invention, can be determined by those of ordinary skill in the art.

The IL-12 of the present invention can be obtained from a suitable source for use in the present method. For  
20 example, IL-12 can be purified from natural sources (e.g., human, animal), produced by chemical synthesis or produced by recombinant DNA techniques as described in Example 1. In addition, the IL-12 of the present invention include  
25 nucleic acid sequences encoding IL-12, as well as the RNAs encoded by such nucleic acid sequences. As used herein, "interleukin-12" and "IL-12" refer to interleukin 12, its individual subunits, fragments thereof which exhibit IL-12  
adjuvant activity and functional equivalents of "interleukin-12" and "IL-2". Functional equivalents of  
30 "interleukin-12" and "IL-12" include modified IL-12 protein such that the resulting IL-12 product has the same adjuvant activity as the IL-12 described herein, and nucleic acid sequences which through the degeneracy of the genetic code  
encode the same peptide gene product as IL-12 and having  
35 the IL-12 adjuvant activity described herein. For example,

-6-

a functional equivalent of "interleukin-12" and "IL-12" can contain a "silent" codon or amino acid substitution (e.g., substitution of one acidic amino acid for another acidic amino acid; or substitution of one codon encoding a hydrophobic amino acid to another codon encoding a hydrophobic amino acid).

IL-12, a heterodimeric cytokine predominantly excreted by the macrophage cells, has been reported to enhance NK cell and CTL activity, to stimulate the differentiation of Th1 cells, and to induce production of cytokines, such as IFN- $\gamma$  (Gately, M.K., et al, *Cell. Immunol.* 143:127 (1992); Naume, B., et al, *J. Immunol.* 148:2429 (1992); Hsieh, C.S., et al *Science* 260:547 (1993); Manetti, R., et al *J. Exp. Med.* 177:1199 (1993); Chan, S.H., et al, *J. Exp. Med.* 173:869 (1991); D'Andrea, A., et al, *J. Exp. Med.* 176:1387 (1992); Macatonia, S.E., et al, *Int. Immunol.* 5:1119 (1993); Tripp, C.S., et al, *Proc. Natl. Acad. Sci. USA* 90:3725 (1993)). IL-12 formerly referred to as natural killer cell stimulatory factor or cytotoxic lymphocyte maturation factor functions to activate and to link the innate and acquired immune responses (Kobayashi, M., et al *J. Exp. Med.* 170:827 (1989); Stern, A. S., et al *Proc. Natl. Acad. Sci. USA* 87:6808 (1990); Locksley, R.M., et al *Proc. Natl. Acad. Sci. USA* 90:5879 (1993)). IL-12 promotes differentiation of uncommitted T helper cells towards the Type 1 (Th1) phenotype (Hsieh, C.S., et al *Science* 260:547 (1993); Manetti, R., et al *J. Exp. Med.* 177:1199 (1993)). This results in a characteristic constellation of cytokines, such as IFN- $\gamma$ , and generally promotes cell-mediated immunity (Chan, S.H., et al, *J. Exp. Med.* 173:869 (1991); D'Andrea, A., et al, *J. Exp. Med.* 176:1387 (1992) Macatonia, S.E., et al, *Int. Immunol.* 5:1119 (1993); Gately, M.K., et al, *Cell. Immunol.* 143:127 (1992) Naume, B., et al, *J. Immunol.* 148:2429 (1992)). IL-12 has been demonstrated to enhance the immune response and



to improve protective immunity in several infectious disease models, including Listeriosis, Leishmaniasis, Toxoplasmosis and lymphocytic choriomeningitis virus infection (Tripp, C.S., et al, *Proc. Natl. Acad. Sci. USA* 90:3725 (1993); Heinzl, F.P., et al *J. Exp. Med.* 177:1505 (1993); Sypek, J.P., et al *J. Exp. Med.* 177:1797 (1993); Afonso, L. C., et al, *Science* 263:235 (1994); Gazzinelli, R.T., et al, *Proc. Natl. Acad. Sci. USA* 90:6115 (1993); Khan, I.A., et al *Infect. Immun.* 62:1639 (1994); Orange, J.S. et al, *J. Immunol.* 152:1253 (1994)). The purification and cloning of IL-12 are disclosed in PCT publication nos. WO 92/05256 and WO 90/05147, and in European patent publication no. 322,827 (identified as "CLMF").

Interleukin-12 or IL-12 is a mammalian cytokine which exhibits numerous immunologic effects, including modulation of T cell response to antigens (see, for example, PCT publication nos. WO 92/05256 and WO 90/05147, wherein IL-12 is identified as "NKSF"). It has also been suggested generally that IL-12 might have some application as a vaccine adjuvant (Scott, P., *Science*, 260:496-497(1993); Trichieri, G., *Immunology Today*, 14:335-338(1993)).

In the method of the present invention, an effective adjuvant amount of IL-12 is administered in combination with an antigen of a virus of the Paramyxoviridae family. That is, the IL-12 is administered at a time closely related to immunization of the host with the viral antigen, so that an enhanced immune response in the host is produced relative to the immunization of a host in which IL-12 is not administered. Thus, the IL-12 can be administered prior to, preferably just prior to, immunization, at the time of immunization (i.e., simultaneously) or after immunization (i.e. subsequently). In addition, the IL-12 can be administered prior to immunization with the viral antigen of the Paramyxoviridae family, followed by

subsequent injections of IL-12 after immunization with the antigen.

The IL-12 and the antigen can be administered to a host in a variety of ways. The routes of administration include intradermal, transdermal (e.g., slow release polymers), intramuscular, intraperitoneal, intravenous, subcutaneous, oral, epidural and intranasal routes. Any other convenient route of administration can be used, for example, infusion or bolus injection, or absorption through epithelial or mucocutaneous linings. In addition, the IL-12 and the antigen of the Paramyxoviridae virus can be administered together with other components or biologically active agents, such as other known adjuvants (e.g., alum, MPL, QS21), pharmaceutically acceptable surfactants (e.g., glycerides), excipients (e.g., lactose), carriers, diluents and vehicles. If desired, certain sweetening, flavoring and/or coloring agents can also be added.

The IL-12 and the antigen can be administered as a prophylactic vaccine to hosts which are either infected or uninfected with the virus. The IL-12 and the antigen can also be administered as a therapeutic vaccine to infected hosts and can result in amelioration or elimination of the disease state caused by the infecting virus.

Further, the antigen and/or IL-12 can be administered by *in vivo* expression of polynucleotides encoding such into a mammalian subject. For example, the IL-12 or the Paramyxoviridae antigen can be administered to a host using live vectors, wherein the live vectors containing IL-12 and/or antigen nucleic acid sequences are administered under conditions in which the antigen and/or IL-12 are expressed in vivo. For example, a host can be injected with a vector which encodes and expresses an antigen of a virus of the Paramyxoviridae family in vivo in combination with IL-12 protein or peptide, or in combination with a vector which encodes and expresses the IL-12 protein in

- vivo. Alternatively, a host can be injected with a vector which encodes and expresses IL-12 in vivo in combination with a Paramyxoviridae antigen in peptide or protein form, or in combination with a vector which encodes and expresses
- 5 a Paramyxoviridae antigen in vivo. A single vector containing the sequences encoding a Paramyxoviridae antigen and the IL-12 protein are also useful in the methods and compositions of the present invention.

- Several expression vector systems are available
- 10 commercially or can be reproduced according to recombinant DNA and cell culture techniques. For example, vector systems such as the yeast or vaccinia virus expression systems, or virus vectors can be used in the methods and compositions of the present invention (Kaufman, R.J., A J.
- 15 of Meth. in Cell and Molec. Biol., 2:221-236 (1990)). Other techniques using naked plasmids or DNA, and cloned genes encapsidated in targeted liposomes or in erythrocytes ghosts, can be used to introduce the IL-12 and/or Paramyxoviridae antigen polynucleotides into the host
- 20 (Freidman, T., Science, 244:1275-1281 (199); Rabinovich, N.R., et al., Science, 265:1401-1404 (1994)). The construction of expression vectors and the transfer of vectors and nucleic acids into various host cells can be accomplished using genetic engineering techniques, as
- 25 described in manuals like *Molecular Cloning* and *Current Protocols in Molecular Biology* or by using commercially available kits (Sambrook, J., et al., *Molecular Cloning*, Cold Spring Harbor Press, 1989; Ausubel, F.M., et al., *Current Protocols in Molecular Biology*, Greene Publishing
- 30 Associates and Wiley-Interscience, 1989).

- The amount of antigen used in the methods and compositions of the present invention is an amount which produces an effective immunostimulatory response in the host. An effective adjuvant amount of IL-12 is an amount
- 35 such that when administered, it results in an enhanced

-10-

immune response relative to the immune response when an effective adjuvant amount of IL-12 is not administered. In addition, the amount of antigen from a virus of the Paramyxoviridae family and IL-12 used to immunize the host will vary depending on a variety of factors, including the antigen employed, the size, age, body weight, general health, sex and diet of the host, and the time of administration, duration or particular qualities of the Paramyxoviridae virus being vaccinated against. Adjustment and manipulation of established dosage ranges are well within the ability of those skilled in the art.

The formulation and route of delivery of vaccine products can influence the induction of T helper lymphocyte subsets and may thereby affect disease expression after viral challenge (Graham, B.S., et al, *Immunol.* 151:2032 (1993)). Depletion of IL-4 at the time of immunization by neutralizing monoclonal antibody induces a Th2 to Th1-like immune response shift, accompanied by an improved clinical outcome and an increased CD8+ cytotoxic T lymphocyte (CTL) activity (Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). This was associated with an increased expression of endogenous IL-12 message at the time of challenge in the anti-IL-4 treated mice. These findings suggest that selective activation of the Th2-like cell subset may be responsible for RSV vaccine induced immunopotentialization of disease and that IL-12 may be associated with shifting the response away from Th2 to a more Th1-like response.

Replication of RSV is markedly reduced after live RSV challenge in mice given IL-12. Use of IL-12 as an adjuvant in a composition comprised of an antigen from RSV and IL-12 also induced a shift from a Th2 to a Th1-like immune response in mice after RSV challenge. While the IL-12 adjuvant effect was potent for reduction of RSV replication, it is more important for use as a vaccine adjuvant to decrease illness following challenge by RSV.

-11-

The use of cytokines as adjuvants can allow one to control the immune parameters induced by immunization to improve protective effects and decrease the negative effects of a vaccine for RSV. The effects of IL-12 on the immune responses to RSV vaccination as measured after live virus challenge in the BALB/c mouse model are described in the Examples. The results indicate that IL-12 acts as a potent adjuvant and is a useful product to include in RSV vaccines.

As described in Example 1, BALB/c mice were immunized with inactivated whole virus intramuscularly, and murine recombinant IL-12 was administered intraperitoneally for 5 successive days starting at one day before immunization or challenge. The mice were challenged with live virus 4 weeks later. The viral replication in lungs 4 days after challenge was assessed. IL-12 administered at the time of immunization had a marked effect on the reduction of viral replication. Log 10 pfu per lung was reduced from 6.9 in RSV-immunized control mice without IL-12 administration to 3.8 in immunized mice with IL-12 administration (Figure 1). In contrast, IL-12 administration at the time of challenge did not have significant effect on the viral replication (Figure 1).

The effects of different delivery routes for IL-12 is described in Example 2. IL-12 was administered either intraperitoneally as described in Example 1, or intramuscularly mixed with the RSV immunogen. Table 1 shows the effect of either intraperitoneal or intramuscular delivery of IL-12 on the reduction of viral replication. A single dosage of IL-12 given simultaneously with immunogen had the same effect on the reduction of viral replication compared to the 5-dosage intraperitoneal regimen. IL-12 as a specific immunomodulator only worked in RSV immunized mice, having no effects on the unprimed mice (Figure 1,

-12-

Table 1). These data demonstrate that IL-12 exerts a potent adjuvant effect on the inactivated RSV immunogen.

The patterns of immunoglobulin isotypes produced in response to immunization are indirect indicators of the types of cytokines produced *in vivo*. IgG2a is produced in mice as a consequence of Th1 cell activation, whereas IL-4 promotes the production of IgG1 (Burststein, H.J. et al, *J. Immunol.* 147:2950 (1991); Finkelman, F.D., et al, *Annu Rev. Immunol.* 8:303 (1990); Morris, S.C., et al, *J. Immunol.* 152:1047 (1994)). As described in Example 3, blinded assays of serum RSV-specific immunoglobulin isotype titers in RSV-immunized mice receiving IL-12 showed that IL-12 induced significantly more RSV-specific IgG2a antibody and significantly less IgG1 antibody compared to immunized mice not given IL-12. The pattern of IgG2a and IgG1 RSV-specific antibody response was similar whether IL-12 was given intramuscularly or intraperitoneally (Table 2).

The pattern of antibody isotype utilization induced by IL-12 suggests that *in vivo* IL-12 administration can promote the differentiation of antigen-specific CD4+ Th1 cells and inhibit the development of Th2 cells in response to the inactivated RSV intramuscular immunization. The pattern of cytokine mRNA expression in lungs was directly examined as described in Example 4. Lung tissues from immunized mice, with or without IL-12 treatment, were harvested at 4 days after live virus challenge. The cytokine mRNAs for IFN- $\gamma$ , IL-4, IL-6, IL-10, and IL-12, were measured by Northern blot analysis. There were no obvious differences in IL-6 and IL-10 mRNA levels among mice with different treatments. However, the lungs from mice treated with IL-12 at the time of either immunization or challenge contained more IFN- $\gamma$  relative to IL-4 compared to control mice that did not receive IL-12 (Figure 2). Increased IL-12 mRNA expression occurred in the mice treated with IL-12 at the time of immunization, while IL-12

-13-

administration at challenge did not increase IL-12 mRNA expression (Figure 2). These data suggest that IL-12 enhanced Th1 cell differentiation and produced a shift from a Th2 to a Th1-like response in mice immunized with the  
5 inactivated RSV immunogen.

RSV-specific cytotoxic T lymphocyte (CTL) activity in lungs of immunized mice was assessed to evaluate whether IL-12 administration enhanced cell-mediated immunity as a positive modulatory effector. As described in Example 5, a  
10 direct CTL assay using lung lymphocytes was employed which does not include *in vitro* stimulation (Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). There was no difference in CTL activity between groups that received or did not receive IL-12 treatment at the time of immunization. This result,  
15 which was repeated in two consecutive experiments, further suggests the Th1-like cytokine pattern was a product of CD4+ T cells. It is reasonable to expect that altering the dose of IL-12 would induce a greater CD8+ response which would alter the illness pattern.

20 As described in Example 5, even though IL-12 has a dramatic effect on the reduction of RSV replication in lungs, there was not significant difference in the clinical outcome, including weight loss and illness score between groups. The simple shift in the pattern of cytokine  
25 expression was therefore not predictive of a change in illness. This suggests that the cell populations responsible for cytokine production can be key determinants of illness and not cytokines themselves. For example, mice treated with anti-IL-4 at the time of immunization also had  
30 increased IFN- $\gamma$  expression in lungs at the time of challenge. However, the anti-IL-4 treatment resulted in diminished illness that was associated with increased CD8+ CTL activity (Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). In the case of anti-IL-4 treated mice, it may be that the

-14-

IFN- $\gamma$  was a product of CD8+ T cells, whereas in IL-12 treated mice, CD4+ T cells are a more likely source.

- Adjusting the dose of IL-12 can alter its properties. A recent study of IL-12 on immune responses to lymphocytic choriomeningitis (LCMV) infection showed that low doses of IL-12 enhanced immunity to LCMV infection as demonstrated by increased splenic CD8+ T cell numbers and decreased LCMV replication. However, high doses of IL-12, equivalent to those used in the examples, impaired resistance against LCMV infection as demonstrated by reduced virus-specific CTL activity and increased viral replication (Orange, J.S. et al, *J. Immunol.* 152:1253 (1994)). It may therefore be possible to adjust the dose of IL-12 or its method of delivery to maintain the effect on viral inhibition, but to also impact illness.

The invention is further illustrated in the following examples.

#### EXEMPLIFICATION

##### Example 1 Immunization of mice with RSV and IL-12

- 20 Mice: Pathogen-free female BALB/c mice, 8 to 10 months old, were purchased from Charles River Laboratories (Raleigh, North Carolina) and cared for according to the "Guide for the Care and Use of Laboratory Animals" as previously described (Graham, B.S., et al, *J. Med. Virol.* 26:153 (1988)).

- RSV Immunogen and Virus: Preparation of the formalin-inactivated alum-precipitated RSV and preparation of stock of the A2 strain of RSV have been previously reported (Graham, B.S., et al, *Immunol.* 151:2032 (1993)).
- 30 Both the vaccine preparation and the challenge stock were derived from the A2 strain of RSV.

Murine Cytokine IL-12: Murine recombinant IL-12 was expressed from cloned cDNAs (Schoenhaut, D.S., et al, *J.*



-15-

- Immunol.* 148:3433 (1992)). The lot used in this paper was MRB021693-1.2 (Genetics Institute, Cambridge, Massachusetts) with a specific activity of  $5.6 \times 10^6$  units/mg as determined by PHA blast assay (Wolf, S.F., et al, *J. Exp. Med.* 146:3074 (1991)). Concentrated aliquots of IL-12 were stored at  $-70^\circ\text{C}$  and diluted in phosphate-buffered saline with 1% normal mouse serum (1% PBS).
- Immunization: Mice were immunized with formalin-inactivated alum-precipitated RSV containing  $2.2 \times 10^6$  pfu equivalents of virus antigen intramuscularly, and challenged with  $10^7$  pfu of live RSV intranasally 4 weeks later as previously described (Graham, B.S., et al, *J. Immunol.* 151:2032 (1993)); Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). IL-12 was administered intraperitoneally for 5 successive days, starting at one day before immunization at a dose of  $1 \mu\text{g}/\text{mouse}$ . Control mice received 1% phosphate buffered saline (PBS) on the same schedule.

- The viral replication in lungs 4 days after challenge was assessed by plaque assay. Mouse serum samples were collected on the day of and two weeks after live RSV challenge.

- Plaque Assays and Neutralization Tests: Two-day old HEp-2 monolayers, 80% confluent in Costar 12-well plates, were used for plaque assay and neutralization tests. The assays were performed as described previously (Graham, B.S., et al, *J. Med. Virol.* 26:153 (1988)).

- IL-12 administered at the time of immunization has a marked effect on the reduction of viral replication. Log 10 pfu per lung was reduced from 6.9 in RSV-immunized control mice without IL-12 administration to 3.8 in immunized mice with IL-12 administration. In contrast, IL-12 administration at the time of challenge did not have significant effect on the viral replication. See Figure 1.

-16-

(Log 10pfu/gram lung is shown as arithmetic means  $\pm$  S.D.;  
KV denotes killed virus)

Example 2 Effect of delivery route of IL-12 on its adjuvant  
ability

5       The effect of a different delivery route of the  
adjuvant IL-12 was assessed. IL-12 was administered  
intraperitoneally as described in Example 1 to one group of  
mice. To another group of mice, IL-12 was administered  
intramuscularly mixed with the RSV antigen. The control  
10 mice were either mock immunized or treated with IL-12 alone  
without antigen. Table 1 summarizes the results of the  
experiment which show that a single dosage of IL-12 given  
simultaneously with immunogen had the same effect on the  
reduction of viral replication compared to the 5-dosage  
15 intraperitoneal regimen. IL-12 as a specific  
immunomodulator only worked in RSV immunized mice, having  
no effects on the unprimed mice. See Figure 1 and Table 1.  
These data demonstrate that IL-12 exerted a potent adjuvant  
effect on the inactivated RSV immunogen.

20 Example 3 Assay of immunoglobulin isotype titers in RSV-  
immunized mice receiving IL-12

The patterns of immunoglobulin isotypes produced in  
RSV-immunized mice receiving IL-12 was examined. Mouse  
serum samples were collected on the day of and two weeks  
25 after live RSV challenge.

RSV-Specific Immunoglobulin Isotype ELISA: All serologic  
assays were performed by a person blinded to the  
experimental groups. BCH4 and BC cells were bound to the  
solid phase on Immulon II 96-well plates (NUNC, Denmark).  
30 Serial diluted mouse serum samples were added to each well.  
Plates were incubated, washed, and goat anti-murine IgG1 or  
IgG2a conjugated to alkaline phosphatase (Southern  
Biotechnology, Birmingham, Alabama) diluted 1:1000 was

-17-

- added, respectively. After another incubation, plates were washed and substrate was added for 30 minutes at room temperature and OD<sub>405</sub> was determined (Graham, B.S., et al, *Immunol.* 151:2032 (1993); Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). A serum dilution was considered positive if the mean optical density of two BCH4 cell wells was greater than twice that of BC-coated wells and greater than 0.1.

- The results demonstrate that IL-12 induced significantly more RSV-specific IgG2a antibody and significantly less IgG1 antibody compared to immunized mice not given IL-12. The pattern of IgG2a and IgG1 RSV-specific antibody response was similar whether IL-12 was given intramuscularly or intraperitoneally. See Table 2.

15 Example 4 Pattern of cytokine mRNA expression in lungs from mice immunized with and without IL-12

- Lung tissues from immunized mice, with and without IL-12 treatment, were harvested at 4 days after live virus challenge. The cytokine mRNAs for IFN- $\gamma$ , IL-4, IL-6, IL-10 and IL-12 were measured by Northern blot analysis.

- mRNA Extraction, Northern Blotting, and Cytokine Detection: The total RNA from whole lungs was extracted and polyA RNA isolated, electrophoretically separated and transferred to membrane as previously described (Graham, B.S., et al, *Immunol.* 151:2032 (1993); Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). Hybridization with <sup>32</sup>P oligonucleotide probes were performed as previously described (Graham, B.S., et al, *Immunol.* 151:2032 (1993)). After washing, membranes were exposed to Kodak X-omat film at -70°C. Laser densitometry was performed with an LKB Ultrosan XL using GelScan XL software (Pharmacia Fine Chemicals, Piscataway, New Jersey). Oligonucleotide probes for murine IL-4, IL-10, IFN- $\gamma$ , IL-6 were purchased from R &

-18-

D Systems (Minneapolis, Minnesota) or Clontech Laboratory Inc. (Palo Alto, California). A cocktail of oligonucleotides designed for detecting IL-12 was based on the murine IL-12 sequence spanning predicted splice sites  
5 based on those identified in the human IL-12 sequence. (Schoenhaut, D.S., et al, *J. Immunol.* 148:3433 (1992)). From 5' to 3':

p-40:

10 TGAGGACACATCTTGCTTTGCTGCGAGCTG (SEQ. ID. NO:1),  
TCCCGCCTTTGCATTGGACTTCGGTGATG (SEQ. ID. NO:2), and  
CAACGTTGCATCCTAGGATCGGACCCTGCA (SEQ. ID. NO:3);

p-35:

GCCAGGCAACTCTCGTTCTTGTGTAGTTCC (SEQ. ID. NO:4), and  
GCGTTGATGGCCTGGAATCTGTCTGGTAC (SEQ. ID. NO:5).

15 Cytokine mRNA Northern blots (2 samples per group) with averaged  $\alpha$ -tubulin-normalized densities are shown in Figure 2. There were no obvious differences in IL-6 and IL-10 mRNA levels among mice with different treatments. However, the lungs from mice treated with IL-12 at the time  
20 of either immunization or challenge contained more IFN- $\gamma$  relative to IL-4 compared to control mice that did not receive IL-12 (Figure 2). An increased IL-12 mRNA expression occurred in the mice treated with IL-12 at the time of immunization, while IL-12 administration at  
25 challenge did not increase IL-12 mRNA expression (Figure 2). These data suggested that IL-12 enhanced Th1 cell differentiation and produced a shift from a Th2 to a Th1-like response in mice immunized with the inactivated RSV immunogen.

30 Example 5 RSV-specific cytotoxic T lymphocyte (CTL)  
activity in lungs of immunized mice

RSV-specific CTL activity in lungs of immunized mice was assessed to evaluate whether IL-12 administration

-19-

enhanced cell-mediated immunity as a positive modulatory effector. A direct CTL assay using lung lymphocytes was employed which does not include *in vitro* stimulation (Tang, Y.-W., et al, *J. Clin. Invest.* (1994)).

- 5 Cytotoxicity T Cell Assays: Whole lung lymphocytes were isolated by Ficoll-Hypaque (1.09 specific gravity) cushion centrifugation. BCH4 and BC target cells labeled with  $^{51}\text{Cr}$  (Dupont-New England Nuclear, Boston, Massachusetts) were incubated with effector cells for 4 hours at 37°C in 96-  
10 well microtiter plates as described (Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). The spontaneous and total release were obtained by treating the target cells with 10% RPMI and 5% Triton X-100 detergent, respectively. Each point was the mean from three replicate wells. The specific  
15 release of  $^{51}\text{Cr}$  from target cells was defined as  $100 \times (\text{sample cpm} - \text{background cpm}) / (\text{total cpm} - \text{background cpm})$ .

- There was no difference in CTL activity between groups that received or did not receive IL-12 treatment at the time of immunization. This result was repeated in two  
20 consecutive experiments and further suggests the Th1-like cytokine pattern was a product of CD4+ T cells. Even though IL-12 has a dramatic effect on the reduction of RSV replication in lungs, there was no significant difference in the clinical outcome, including weight loss and illness  
25 score between groups. Illness assessment, including weight loss and clinical scores were performed as previously described (Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). The simple shift in the pattern of cytokine expression was therefore not predictive of a change in illness. This  
30 suggests that the cell population responsible for cytokine production may be a key determinant of illness and not cytokines themselves. For example, mice treated with anti-IL-4 at the time of immunization also had increased IFN- $\gamma$

-20-

expression in lungs at the time of challenge. However, the anti-IL-4 treatment resulted in diminished illness that was associated with increased CD8+ CTL activity (Tang, Y.-W., et al, *J. Clin. Invest.* (1994)). In the case of anti-IL-4  
5 treated mice, it may be that the IFN- $\gamma$  was a product of CD8+ T cells, whereas in IL-12 treated mice, CD4+ T cells were a more likely source.

#### EQUIVALENTS

Those skilled in the art will recognize, or be able to  
10 ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described specifically herein. Such equivalents are intended to be encompassed in the scope of the following claims.

-21-

TABLE 1

IL-12 Administered Either Intraperitoneally or  
Intramuscularly Reduces Virus Replication  
in Lungs and Noses

Group	N	Immunogen	IL-12 Route	Log 10pfu/gram Lung*	Log 10pfu/ Nose*
1	5	KV	--	6.56 ± 0.44	3.73 ± 0.15
2	5	KV	IP	4.75 ± 0.25†	2.61 ± 0.23‡
3	5	KV	IM	4.44 ± 0.55†	2.80 ± 0.34‡
4	5	Mock	--	6.55 ± 0.32	3.73 ± 0.29
5	5	Mock	IP	6.51 ± 0.34	3.21 ± 0.27

\* Mean ± S.D.

† p < 0.001 compared to Log 10pfu/gram lung in group 1

‡ p < 0.001 compared to log 10pfu/nose in group 1

KV killed virus

IP intraperitoneal

IM intramuscularly

TABLE 2

IL-12 Administered Alters RSV-Specific Serum Immunoglobulin Isotype Titers  
at and Two Weeks After RSV Challenge

Group	Immunogen	IL-12 Route	At Challenge		2 Weeks After Challenge	
			IgG1	IgG2a	IgG1	IgG2a
1	KV	--	<80 (0/5)*	<80 (0/5)	640.0 $\pm$ 2.2 (4/4)	269.1 $\pm$ 4.2 (2.4)
2	KV	IP	<80 (0/5)	557.2 $\pm$ 6.4† (3/5)	121.3 $\pm$ 1.9‡ (2/5)	640.0 $\pm$ 6.7‡ (3/5)
3	KV	IM	<80 (0/5)	367.6 $\pm$ 4.1 (3/5)	<80 (0/5)	905.1 $\pm$ 4.2‡ (3.4)
4	Mock	--	<80 (0/5)	<80 (0/5)	<80 (0/5)	113.1 $\pm$ 2.1 (1/4)
5	Mock	IP	<80 (0/5)	<80 (0/5)	<80 (0/5)	139.3 $\pm$ 2.5 (2/5)

\* Number converted/number tested

† Geometric mean titer  $\pm$  S.D. Negative samples were assigned a titer value of 80 for statistical calculations.

‡ p < 0.01 compared to IgG1 2 weeks after challenge in group 1

‡‡ p > 0.5 compared to IgG2a 2 weeks after challenge in group 1

KV killed virus

IP intraperitoneal

IM intramuscular



-23-

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: Vanderbilt University  
(B) STREET: 405 Kirkland Hall  
(C) CITY: Nashville  
(D) STATE/PROVINCE: TN  
(E) COUNTRY: US  
(F) POSTAL CODE/ZIP: 37240  
(G) TELEPHONE: (615) 322-8333  
(I) TELEFAX: (615) 343-3930

(ii) TITLE OF INVENTION: INTERLEUKIN-12 AS AN ADJUVANT FOR  
PARAMYXOVIRIDAE VACCINES

(iii) NUMBER OF SEQUENCES: 5

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Hamilton, Brook, Smith & Reynolds, P.C.  
(B) STREET: Two Militia Drive  
(C) CITY: Lexington  
(D) STATE: MA  
(E) COUNTRY: US  
(F) ZIP: 02173

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/318,480  
(B) FILING DATE: October 5, 1994  
(C) CLASSIFICATION:

-24-

## (viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Brook Esq., David E.
- (B) REGISTRATION NUMBER: 22,592
- (C) REFERENCE/DOCKET NUMBER: VDB94-01 PCT

## (ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (617) 861-6240
- (B) TELEFAX: (617) 861-9540

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TGAGGACACA TCTTGCTTTG CTGCGAGCTG

30

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCCCGCCTTT GCATTGGACT TCGGTGATG

29

-25-

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CAACGTTGCA TCCTAGGATC GGACCCTGCA

30

## (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GCCAGGCAAC TCTCGTTCTT GTGTAGTTCC

30

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 30 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCGTTGATGG CCTGGAAGTC TGCTGGTAC

30

-26-

CLAIMS

We claim:

1. A composition comprising a mixture of an antigen of a virus of the Paramyxoviridae family and an effective  
5 adjuvant amount of interleukin-12.
2. A composition of Claim 19 wherein the virus of the Paramyxoviridae family is selected from the group consisting of: parainfluenza virus 1, parainfluenza virus 2, parainfluenza virus 3, parainfluenza virus 4,  
10 mumps virus, measles virus, respiratory syncytial virus, canine distemper virus, bovine respiratory syncytial virus, Newcastle disease virus and rhinderpest virus.
3. A composition of Claim 19 wherein the antigen is  
15 selected from the group consisting of: the fusion glycoprotein, the hemagglutinin-neuraminidase and the G glycoprotein.
4. A composition comprising a mixture of an antigen of a virus of the Paramyxoviridae family and an adjuvant  
20 amount of interleukin-12 for use in reducing viral replication of virus of the Paramyxoviridae family in a host.
5. A composition comprising a mixture of an antigen of a virus of the Paramyxoviridae family and an adjuvant  
25 amount of interleukin-12 for use in eliciting an immune response against a virus of the Paramyxoviridae family in a host.

-27-

6. A composition according to Claim 4 or 5 wherein the virus of the Paramyxoviridae family is selected from the group consisting of: parainfluenza virus 1, parainfluenza virus 2, parainfluenza virus 3,  
5 parainfluenza virus 4, mumps virus, measles virus, respiratory syncytial virus, canine distemper virus, bovine respiratory syncytial virus, Newcastle disease virus and rhinderpest virus.
7. A composition of Claim 6 wherein the antigen is  
10 selected from the group consisting of: the fusion glycoprotein, the hemagglutinin-neuraminidase and the G glycoprotein.
8. A composition according to Claim 4 or 5 wherein the antigen and/or interleukin-12 is a polynucleotide  
15 expressed in the host.
9. A method of reducing viral replication of a virus of the Paramyxoviridae family in a host, comprising administering to the host an antigen of the virus in combination with an effective adjuvant amount of  
20 interleukin-12.
10. A method of eliciting an immune response against a virus of the Paramyxoviridae family in a host, comprising administering to the host an antigen of the virus in combination with an effective adjuvant amount  
25 of IL-12.

1/1




FIG. 1




FIG. 2

## INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US 95/12656A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K39/155 A61K39/39

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SCIENCE (WASHINGTON D C) 263 (5144). 1994. 235-237, AFONSO L C C ET AL 'The adjuvant effect of interleukin - 12 in a vaccine against Leishmania major.' cited in the application see the whole document ---	1-10
Y	IMMUNOLOGY TODAY, vol. 14, no. 7, July 1993 CAMBRIDGE GB, pages 335-337, TRINCHIERI G. 'Interleukin-12 and its role in the generation of TH1 cells' cited in the application see the whole document ---	1-10
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- 'A' document defining the general state of the art which is not considered to be of particular relevance  
'E' earlier document but published on or after the international filing date  
'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
'O' document referring to an oral disclosure, use, exhibition or other means  
'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
'A' document member of the same patent family

Date of the actual completion of the international search

20 February 1996

Date of mailing of the international search report

19.03.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentian 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040; Tx 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Moreau, J

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US 95/12656

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JOURNAL OF IMMUNOLOGY, vol. 151, no. 4, 15 August 1993 BALTIMORE US, pages 2032-2040, GRAHAM B.S. ET AL. 'Priming Immunization Determines T Helper Cytokines mRNA Expression Patterns in Lungs of Mice Challenged with Respiratory Syncytial Virus' cited in the application see the whole document ---	1-10
Y	JOURNAL OF IMMUNOLOGY, vol. 152, no. 3, 1 February 1994 BALTIMORE US, pages 1253-1264, ORANGE J.S. ET AL. 'Effects of IL-12 on the Response and Susceptibility to Experimental Viral Infections' cited in the application see the whole document ---	1-10
Y	SCIENCE (WASHINGTON D.C.), vol. 265, no. 5177, 2 September 1994 pages 1401-1404, RABINOVITCH N.R. ET AL. 'Vaccine technologies: View to the Future' cited in the application see the whole document ---	1-10
P,Y	WO,A,94 27636 (AMERICAN CYANAMID COMPANY) 8 December 1994 see the whole document ---	1-10
P,X	JOURNAL OF INFECTIOUS DISEASES 172 (3). 1995. 734-738. , TANG Y-W ET AL. 'Interleukin - 12 treatment during immunization elicits a T helper cell type 1-like immune response in mice challenged with respiratory syncytial virus and improves vaccine immunogenicity.' see the whole document ---	1-10
E	WO,A,95 30437 (AMERICAN HOME PRODUCTS CORPORATION) 16 November 1995 see page 5, line 32 see claim 1 -----	1-10



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/12656

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 9-10  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 9-10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims, it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International Application No

PCT/US 95/12656

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9427636	08-12-94	AU-B- 6957194	20-12-94
WO-A-9530437	16-11-95	AU-B- 2438795	29-11-95